

24503093000



LANE MEDICAL LIBRARY STAMFORD
D1144C7 L5 1931
Some experiments on the influence of lig
OLD

OME EXPERIMENTS ON THE INFLUENCE
OF LIGHT AND POISONS ON MARINE
COPEPODS, WITH REFERENCE
TO THEIR DAILY MIGRATION

BY

W. W. LEPESCHKIN

(Contribution from the Laboratories of the Desert Sanatorium
and Institute of Research, Tucson, Arizona)

BULLETIN OF THE SCRIPPS INSTITUTION OF OCEANOGRAPHY
LA JOLLA, CALIFORNIA

TECHNICAL SERIES, Volume 3, No. 2, pp. 33-35

UNIVERSITY OF CALIFORNIA PRESS
BERKELEY, CALIFORNIA

1931

D1144
C7L5
1931

LANE

MEDICAL



LIBRARY

LEVI COOPER LANE FUND

SOME EXPERIMENTS ON THE INFLUENCE
OF LIGHT AND POISONS ON MARINE
COPEPODS, WITH REFERENCE TO
THEIR DAILY MIGRATION

BY

W. W. LEPESCHKIN
W.P.

LANE LIBRARY

BULLETIN OF THE SCRIPPS INSTITUTION OF OCEANOGRAPHY
OF THE UNIVERSITY OF CALIFORNIA
LA JOLLA, CALIFORNIA

TECHNICAL SERIES, Volume 3, No. 2, pp. 33-35

Issued September 30, 1931

UNIVERSITY OF CALIFORNIA PRESS
BERKELEY, CALIFORNIA

CAMBRIDGE UNIVERSITY PRESS
LONDON, ENGLAND

UNIVERSITY OF CALIFORNIA

1931

SOME EXPERIMENTS ON THE INFLUENCE OF LIGHT AND POISONS ON MARINE COPEPODS, WITH REFERENCE TO THEIR DAILY MIGRATION

BY
W. W. LEPESCHKIN

(Contribution from the Laboratories of the Desert Sanatorium and
Institute of Research, Tucson, Arizona)

The daily migration of marine copepods from upper to lower layers of sea water is usually ascribed to their phototactic sensitivity. Some copepods are considered as positively, others as negatively phototactic. It has been pointed out that the kind of their phototaxis may change under laboratory conditions, and it seems very probable that some unknown unfavorable factor under laboratory conditions produces this change.

During my work at the Scripps Institution of Oceanography in June and July, 1930, I made some experiments on the influence of sunlight on copepods. The copepods used were identified by Professor Charles B. Wilson, of the Department of Education of Massachusetts, and I express to him my hearty thanks for his aid.

The copepods taken from the plankton in the morning, with *Paracalanus parvus* prevailing, showed a very marked positive phototaxis in the laboratory room. If they were exposed in test tubes to direct sunlight, however, they migrated in some minutes to the side of the tube opposite the sun, thus showing accordingly a negative phototaxis. The same change in phototaxis was shown by *Oithona similis* and *Corycaeus venustus*, which sometimes prevail in the morning plankton. The copepods occurring at the sea surface at night (for instance at 9:00 P.M.), at the end of June, among which were *Acartia clausii* and *A. tonsa*, *Oithona brevicornis* and *O. nana*, *Corycaeus rostratus*, *Oncaea venusta*, and *Eutерpe acutifrons*, proved to be negatively phototactic even in the laboratory room, and they were all the more negatively phototactic when they were exposed to direct sunlight on the next morning, provided they had been kept in constantly running sea water (the outflow tube was provided with a small funnel covered by bolting cloth).

94319

LANE LIBRARY

It is evident from the above experiments that the intensity of light plays an important part in the movement of copepods. It may be supposed also that direct sunlight is injurious to all kinds of copepods and causes them to swim to a safer place. Although the copepods represent complex organisms with well developed nervous processes, in their phototaxis they are suggestively similar to swimming algae which change their positive to negative phototaxis when the intensity of light is too strong.

That direct sunlight is injurious to copepods is shown by the following experiments. The exposure of various kinds of copepods to direct sunlight decreases their resistance against poisons, for instance, against mercury bichloride.

Sea water containing copepods (the experiments were made on *Paracalanus parvus*, *Oncaea venusta*, and *Eutерpe acutifrons*) was poured into the test tubes. There were twelve tubes and each tube contained about thirty animals in 8 cc of sea water. Half of the tubes were wrapped in black sateen (three layers), all tubes were immersed in running water of an approximately constant temperature (22° C), and then exposed to direct sunlight at noon. After two hours 0.1 cc of differently concentrated solutions of mercury bichloride was added to each tube, and the time necessary to stop the movement of all animals was measured with a stop watch. The stoppage of movement was irreversible and may be considered as death. The experiment was repeated six times and the variation of the time for different tubes with the same concentration of mercury bichloride did not exceed 10 per cent of the average time given below. The animals which had been in the dark showed the following average duration of life in the solutions of mercury bichloride:

Concentration of HgCl ₂ in per cent.....	0.025	0.0125	0.006	0.003	0.0013	0.00026
Survival time in minutes	1.7	2.3	3.8	6.4	10.8	41

The dependence of the life duration upon the concentration of the poison may be expressed approximately by the formula: $\frac{l}{t} = KC^p$ where t is the survival time, C is the concentration, and K and p are constants. The corresponding curve is a so-called adsorption curve. In our case $K = 11.4$ and $p = 0.74$. The copepods which had been exposed to direct sunlight showed a curve of the same form but it was nearer the axis of concentration; in other words the time necessary to kill the animals was shortened, as is seen below:

Concentration of HgCl ₂ in per cent	0.003	0.0013	0.00026
Survival time in minutes	4.5	7.5	30

Thus the exposure to direct sunlight decreases the resistance of copepods to poisons, and is therefore injurious to them. This conclusion concerns, of course, only visible rays of sunlight. The ultraviolet rays of the solar spectrum proved to increase the resistance of the animals to poisons if the exposure to the rays was short. The visible part of sunlight was absorbed in these experiments by the filter 586-A (purple, ultraviolet) produced by the Corning Glass Company. This filter was used as a cover for one of the Petri dishes in which the animals were swimming in sea water. The resistance of the animals was increased by 15 per cent if the exposure lasted not longer than ten minutes. If the exposure was more than one hour the resistance decreased, however. It seems that both effects of ultraviolet rays (harmful and useful) are independent of each other. Indeed, if the dish with the copepods was covered by a quartz plate or open and exposed to direct sunlight, a smaller decrease of resistance was observed than in the case where the copepods had been covered by a glass; sometimes no decrease of resistance could be discovered in the first case even if the exposure lasted more than one hour. It may therefore be concluded that ultraviolet rays protect the animals from the injurious effect of visible light.

In nature, both parts of sunlight, visible and ultraviolet, permeate the upper layers of sea water, but ultraviolet rays of short wave length (which are the most active in protecting copepods) are absorbed to a greater degree by sea water than are the more harmful visible rays (blue and violet). Thus the protective effect of ultraviolet rays is more effective on the sea surface. It is therefore possible that the copepods which are less sensitive to light and show a negative phototaxis only in case of an illumination by direct sunlight come to the sea surface because they are more protected there from the injurious effect of the visible rays of direct sunlight.



Digitized by the Internet Archive
in 2026

https://archive.org/details/IA40153516_0108

BULLETIN OF THE SCRIPPS INSTITUTION OF OCEANOGRAPHY
TECHNICAL SERIES—(Continued)

VOLUME 2. 1929—

1. Methods of Estimating the Significance of Differences in or Probabilities of Fluctuations Due to Random Sampling. Numerical Illustrations, Formulae, and Tables Prepared Especially for Estimating Probabilities when the Usual "Method of Least Squares" or "Normal Law of Error" May Not Apply, by George Francis McEwen. March 1929	1.70
2. Surface Catches of Marine Diatoms and Dinoflagellates made by U.S.S. "Pioneer" in Alaskan Waters in 1924, by Winfred Emory Allen. June 1929	.25
3. The Food Habits of the California Sardine in Relation to the Seasonal Distribution of Microplankton, by Ralph C. Lewis. June 1929	.30
4. Quantitative Studies on Inshore Diatoms and Dinoflagellates of the California Coast in 1925, by Horace M. Buley.	
5. Quantitative Studies on Inshore Marine Diatoms and Dinoflagellates Taken at Five Stations on the East Pacific Coast in 1926, by Dorothea Fox.	
Nos. 4 and 5 in one cover. December 1929	.25
6. A Mathematical Theory of the Vertical Distribution of Temperature and Salinity in Water under the Action of Radiation, Conduction, Evaporation, and Mixing Due to the Resulting Convection. Derivation of a General Theory, and Illustrative Numerical Applications to a Tank, a Lake, and a Region of the North Pacific Ocean, by George Francis McEwen. September 1929	1.25
7. Influence of Varying Oxygen Tensions upon the Rate of Oxygen Consumption of Fishes, by Ansel B. Keys. January 1930	.25
8. Methods in Quantitative Research on Marine Microplankton, by Winfred Emory Allen. January 1930	.25
9. Ecologic Relations of Some Foraminifera, by Richard D. Norton. May 1930	.60
10. Quantitative Studies of Surface Catches of Marine Diatoms and Dinoflagellates taken in Alaskan Waters by the International Fisheries Commission in the Fall and Winter of 1927-1928 and 1929, by Winfred Emory Allen. October 1930	.25
11. Results of Evaporation Studies Conducted at the Scripps Institution of Oceanography and the California Institute of Technology, by George F. McEwen. November 1930	.25
12. A Study of the Selective Action of Decreased Salinity and of Asphyxiation on the Pacific Killifish, <i>Fundulus parvipinnis</i> , by Ansel B. Keys. March 1931	1.00
Index.	

VOLUME 3. 1931—

1. Descriptions of Deep-Sea Bottom Samples from the Western North Atlantic and the Caribbean Sea, by E. M. Thorp. May 1931	.40
2. Some experiments on the Influence of Light and Poisons on Marine Copepods, with Reference to their Daily Migration, by W. W. Lepeschkin. September 1931	.25

LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on
or before the date last stamped below.

--	--	--



